

Coexisting orders in the quarter-filled Hubbard chain with elastic deformations

H.D. Rosales¹ and D.C. Cabra^{1,2,3}

¹*Departamento de Física, Universidad Nacional de la Plata, C.C. 67, (1900) La Plata, Argentina*

²*Institut de Physique et Chimie des Matériaux de Strasbourg, UMR 7504, CNRS-UdS, 23 rue du Loess, BP 43, 67034 Strasbourg Cedex 2, France*

³*Facultad de Ingeniería, Universidad Nacional de Lomas de Zamora, Cno. de Cintura y Juan XXIII, (1832) Lomas de Zamora, Argentina.*

(Dated: July 16, 2009)

The electronic properties of the quarter-filled extended Peierls-Holstein-Hubbard model that includes lattice distortions and molecular deformations are investigated theoretically using the bosonization approach. We predict the existence of a wide variety of charge-elastic phases depending of the values of the Peierls and Holstein couplings. We include the effect of the Peierls deformation in the nearest-neighbor repulsion V , that may be present in real materials where Coulomb interactions depend strongly on the distance, and we show that the phase diagram changes substantially for large V when this term is taken into account.

PACS numbers: 71.30.+h, 71.45.Lr, 75.30.Fv, 74.70.Kn

I. INTRODUCTION

The electronic properties of the low dimensional interacting electron systems have attracted great interest for many different reasons, the main being that they are simpler to analyze than higher dimensional ones and they could then be used to gain some insight on their higher dimensional counterparts. Typically this connection can be envisaged by coupling many 1D systems to build up a higher dimensional array. They are also interesting on their own since many materials show a quasi-one-dimensional behavior in a certain range of temperatures where higher dimensional couplings can be neglected. A third motivation to study 1D systems comes from the very exciting cold atom systems which provide a test laboratory for many important theoretical developments already done and also motivates further analysis that could be in principle directly tested in real systems.

In particular, and in connection with real materials, quarter-filled systems have received a lot of attention since they are very good candidates to describe the so called organics charge transfer salts like the Bechgaard salts $(\text{TMTSF})_2\text{X}$ or $(\text{TMTTF})_2\text{X}$ where $\text{X}=\text{PF}_6, \text{AsF}_6$ ¹. The Hubbard model is the basic model for understanding the electronic properties of these quasi 1-D organic conductors and in some cases, lattice deformations could play a central role. In addition to this, in crystalline materials where the building block of the crystal structure is a large molecule, the vibrational properties of the molecules often have large effects on the electronic properties of the material¹. These materials exhibit a variety of electronic states with spatially inhomogeneous charge, spin and lattice structures.

Several experimental studies have shown clear evidence for the existence of charge order^{2,3,4,5,6,7,8,9,10} in organic charge transfer solids and this has stimulated considerable theoretical efforts^{11,12,13,14,15,16,17,18}. These materials involve Coulomb repulsion, both on-site and between nearest-neighbors (U , V respectively) and are $1/4$ -

filled. Due to the nearest-neighbors Coulomb repulsion, it is expected that in these materials a charge ordering $\bullet - \circ - \bullet - \circ$ sets in, where $\bullet$ and $\circ$ correspond to occupied and unoccupied sites. The charge order $\bullet - \circ - \bullet - \circ$ corresponds to the so called $4k_F$ Charge-Density-Wave (CDW), where $k_F = \pi/4a$ with a the lattice constant. However, for large values of V , and when electron-phonon interactions are included, the ground state corresponds to an ordering like $\bullet - \bullet - \circ - \circ$ ^{11,12,13,14,15,16}. The charge order corresponds to a tetramerized $2k_F$ CDW and it coexists with the $2k_F$ periodic modulation of the intersite distances, called Bond-Ordered-Wave (BOW) or a mixed $2k_F + 4k_F$ BOW. This charge order is called Bond-Charge Density Wave (BCDW). On the other hand, if the Coulomb interaction between two sites r_i and r_j , is exponentially short ranged, *i.e.* of the form $\exp[-\alpha|r_i - r_j|]/r^\xi$ with $\alpha, \xi > 0$, lattice distortions may have an important effect on the Coulomb interactions between nearest-neighbor.

Motivated by the previous discussion we study, using the Abelian bosonization approach and a semiclassical analysis of the effective theory, the ground state phase diagram of the Peierls-Holstein-Hubbard model including the effects of the Peierls deformation in the nearest-neighbor Coulomb interaction V . We show that the ground state at zero temperature indeed favors several period two and period four distortion patterns, stemming from a competition between elastic and charge-spin energy. These patterns spontaneously break translational symmetry, with different phases depending on the Coulomb interaction strengths U and V and on the value of the spin-phonon couplings. A rich phase diagram is obtained, including all the combinations of Charge Order and $2k_F - 4k_F$ deformations as a function of the spin-phonon couplings.

When the effect of the Peierls deformation in the nearest-neighbor Coulomb interaction is not considered, our analysis reproduces previous results obtained with numerical methods¹⁹. The main novelty in our treat-

ment is the inclusion of a nearest neighbors Coulomb repulsion V which depends on the distance between adjacent ions and hence depends on the lattice distortions. The phase diagram then changes substantially indicating that this variation on V should not be neglected in comparing to experiments. Our analysis allows for a description only at a qualitative level, but it has the advantage of being easily generalizable to include other perturbations as longer range Coulomb interactions, to treat arbitrary filling fractions, to add an external magnetic field, etc.

The paper is organized as follows. In Section II we present the model and its analytical treatment. The phonon sector is described in the adiabatic approximation by classical static deformations and we analyze the possible patterns. In Section III, we use the bosonization approach to represent the electrons operators in terms of bosons and analyze the effective description by considering all the relevant perturbation terms as semiclassical potentials. We draw a qualitative phase diagram with from analysis. Special emphasis is put on the characterization of the ground state phases that result from the combination of charge-spin degrees of freedom and elastic effects in different parameter ranges. Finally, in Section IV we present a summary and conclusions of the present work.

II. MODEL HAMILTONIAN

We consider a 1D extended Hubbard model with Hamiltonian given by

$$\begin{aligned}
 H = & - \sum_{j,\sigma} t_{j,j+1} [c_{j,\sigma}^\dagger c_{j+1,\sigma} + c_{j+1,\sigma}^\dagger c_{j,\sigma}] + \\
 & + U \sum_j n_{j,\uparrow} n_{j,\downarrow} + \sum_j V_{j,j+1} n_j n_{j+1} + \\
 & + \sum_j \beta_j n_j
 \end{aligned} \quad (1)$$

where $c_{j,\sigma}$ denotes the annihilation operator of an electron with spin $\sigma(=\uparrow, \downarrow)$ at the j th site, and $n_j = n_{j,\uparrow} + n_{j,\downarrow}$ is the charge density with $n_{j,\sigma} = c_{j,\sigma}^\dagger c_{j,\sigma}$. $t_{i,i+1}$ is the hopping integral, the parameters $U(>0)$ and $V_{j,j+1}(>0)$ denote the magnitudes of the on-site and nearest neighbor interactions and β_j represents the internal molecular deformation. We focus on the $\frac{1}{4}$ -filled case, with the average number of electrons per site $\langle n_i \rangle = 1/2$.

The interaction of the charge degrees of freedom in a homogeneous Hubbard chain ($t_{j,j+1} = t$) with phonons is usually modeled by a linear expansion of the exchange couplings around the non distorted values t ,

$$t_{j,j+1} \simeq t(1 - g_P \delta_{P,j})$$

where $\delta_{P,j}$ is the local lattice distortion and g_P is the electron-phonon interaction. On the other hand, local deformations of the molecules can produce changes of the

on-site (or molecular) orbital energies and can simply be taken into account by a Holstein term

$$g_H \sum_j \delta_{H,j} n_j, \quad (2)$$

where g_H is the electron-phonon coupling constant of the on-site type (in Eq. (1) we change $\beta_j \rightarrow g_H \delta_{H,j}$).

In addition to the purely electronic terms in Eq. (1) and unlike previous analysis, we include the effects of the Peierls deformation in the nearest-neighbor Coulomb interaction V that may be important when the Coulomb interaction between two sites r_i and r_j is exponentially short ranged, *i.e.* $\exp[-\alpha|r_i - r_j|]/r^\xi$. Specifically, we perform a linear expansion of the $V_{j,j+1}$ couplings around the non distorted values V ,

$$V_{j,j+1} \simeq V(1 - g_V \delta_{P,j})$$

where g_V is the coupling constant which measures the effects of the bond deformation on the Coulomb repulsion. The complete Hamiltonian, including the elastic energy in the adiabatic approximation, is written as

$$\begin{aligned}
 H = & -t \sum_{j,\sigma} (1 - g_P \delta_{P,j}) [c_{j,\sigma}^\dagger c_{j+1,\sigma} + c_{j+1,\sigma}^\dagger c_{j,\sigma}] + \\
 & + U \sum_j n_{j,\uparrow} n_{j,\downarrow} + V \sum_j (1 - g_V \delta_{P,j}) n_j n_{j+1} + \\
 & + g_H \sum_j \delta_{H,j} n_j \\
 & + \frac{K_P}{2} \sum_j \delta_{P,j}^2 + \frac{K_H}{2} \sum_j \delta_{H,j}^2
 \end{aligned} \quad (3)$$

where the parameters K_H and K_P are the conventional elastic constants for the on site deformation and the lattice distortion, respectively.

Now, we consider the lattice deformations. At quarter-filled ($\langle n_i \rangle = 1/2$) the most general period four lattice deformations without collective displacement, can be parametrized as

$$\begin{aligned}
 \delta_{P,j} &= \delta_{P,d} \cos(\pi x_j) + \delta_{P,t} \cos(\pi/2 x_j + \xi - \pi/4) \\
 \delta_{H,j} &= \delta_{H,d} \cos(\pi x_j) + \delta_{H,t} \cos(\pi/2 x_j - \zeta)
 \end{aligned} \quad (4)$$

where x_j is the position of the j -th site (lattice constant a has been set to one), and the phases ξ and ζ determine the spatial patterns of tetramerization $\delta_{P,t}$ and $\delta_{H,t}$ respectively, while $\delta_{P,d}$ and $\delta_{H,d}$ are the amplitude of the dimerization in the lattice distortion and the intra-site electron-phonon deformation. This parametrization is the most general supported by bosonization, as such a deformation is commensurate. Period four deformations cause commensurability of relevant perturbations at $\langle n_i \rangle = 1/2$ and provide a mechanism for a charge gap in this regime²⁰. Numerical evidence of the dominance of period four lattice deformations has been obtained from self consistent computations¹⁹. A uniform deformation, leading to global size change, can also appear; this would

produce a uniform shift in t and V , which is inessential to our present analysis.

Before analyzing the case $g_V > 0$, we summarize all previous numerical results in the $1/4$ filled case with $g_V = 0$. When $g_P = g_H = 0$ and $U \rightarrow \infty$ there exists a critical value of V ($V_c = 2t$) for the appearance of a $4k_F$ Charge-Density-Wave if $V > V_c$ Fig. 1(c). For $V < V_c$ the extended Hubbard model corresponds to a Luttinger liquid (LL) with no charge order. For finite U , the value of V_c has been calculated with strong coupling perturbation theory²² and numerical methods²³.

For small positive g_P and g_H , and $V < V_c$, the ground state has a dimerized $4k_F$ Bond-Order-Wave (BOW) with uniform site charges Fig. 1(a). If g_P is sufficiently large (g_H fixed), there is a superposition of the $2k_F$ and $4k_F$ BOWs accompanied by the $\bullet - \bullet - o - o$ charge order. This state corresponds to the Bond-Charge-Density-Wave state (BCDW) Fig. 1(b). On the other hand, at large g_H and small g_P , a $4k_F$ CDW is found, with uniform bonds Fig. 1(c). If we increase g_P , the ground state becomes the $4k_F$ CDW- $2k_F$ SP, where the charge order $\bullet - \bullet - o - o$, is accompanied by a $2k_F$ lattice distortion Fig. 1(d). For $V > V_c$, the $4k_F$ BOW phase is diminished due to the strong Coulomb repulsion V and phases with charge-order dominate the phase diagram. In this work we present new results for a $1/4$ filled band including the electron-phonon interaction (Peierls and Holstein couplings) and lattice deformations in the nearest-neighbor Coulomb interaction V that has not previously been studied. We show that this term may change substantially the phase diagram for large V .

III. BOSONIZATION APPROACH AND SEMICLASSICAL ANALYSIS

In order to analyze semi-quantitatively the low energy properties of the model given by Eq. (3), we use the Abelian bosonization method which is generally powerful for the description of one-dimensional chains (for further details see Ref.^{20,21,24,25,26}). Dimensionless parameters, are used below. They are introduced using t as the energy scale as follows: $g_P \rightarrow \tilde{g}_P = (t/K_P)^{1/2} g_P$, $g_H \rightarrow \tilde{g}_H = (t/K_H)^{1/2} g_H$, $g_V \rightarrow \tilde{g}_V = (t/K_P)^{1/2} g_V$, $\delta_{P,s} \rightarrow \tilde{\delta}_{P,s} = (K_P/t)^{1/2} \delta_{P,s}$ and $\delta_{H,s} \rightarrow \tilde{\delta}_{H,s} = (K_H/t)^{1/2} \delta_{H,s}$ ($s = d, t$). To obtain the corresponding low-energy, we write the fermion operator as²⁰

$$\begin{aligned} c_{x,\sigma} \rightarrow \psi_\sigma(x) &\sim e^{ik_{F,\sigma}x} \psi_{L,\sigma}(x) + e^{-ik_{F,\sigma}x} \psi_{R,\sigma}(x) + \dots \\ &= e^{ik_{F,\sigma}x} e^{-i\sqrt{4\pi}\phi_{L,\sigma}(x)} \\ &\quad + e^{-ik_{F,\sigma}x} e^{i\sqrt{4\pi}\phi_{R,\sigma}(x)} + \dots, \end{aligned} \quad (5)$$

where $k_{F,\sigma}$ are the Fermi momenta for up and down spin electrons and $\phi_{R,L,\sigma}$ are the chiral components of two bosonic fields, introduced as usual in order to bosonize the spin up and down chiral fermion operators $\psi_{R,L,\sigma}$. The dots stand for higher order terms²⁰. They take into account the corrections arising from the curvature of




FIG. 1: Possible lattice deformations and charge density wave ground state in the Peierls-Holstein-Hubbard model at quarter filling. The Grey, black and white circles correspond to site charges of $\langle n \rangle = 0.5, 1$ and 0 , respectively. We reproduce the previous results of¹⁹ and we show two new phases (e), (f) that appear when $g_V > 0$. In this figure (a) The $4k_F$ BOW state, with dimerized bond orders and uniform site charge densities, in (b) represent the BCDW state, with bond orders $2k_F > 4k_F$ and charge order $\bullet - \bullet - o - o$. (c) The $4k_F$ CDW with $\bullet - o - \bullet - o$ charge order and uniform bond order. (d) The $4k_F$ CDW- $2k_F$ SP state, with $\bullet - o - \bullet - o$ charge order and bond orders $2k_F$. For $g_V = 0.2$, we have phase (e) *i.e.*, $4k_F$ CDW- $4k_F$ SP with $\bullet - o - \bullet - o$ charge order together with dimerization bond order. Finally, in the phase (f) ($4k_F$ CDW- $4k_F$ SP + $2k_F$ CDW- $2k_F$ SP) is characterized by three different site charges $\frac{1}{2} - \epsilon'$, $\frac{1}{2} + \epsilon$, $\frac{1}{2} - \epsilon'$, $\frac{1}{2} + \epsilon''$ with bond orders $2k_F > 4k_F$.

the dispersion relation due to the Coulomb interaction. For non-zero Hubbard repulsion U and V , the low energy effective Hamiltonian corresponding to (3) written in terms of the bosonic fields $\phi_\uparrow$ and $\phi_\downarrow$ has a complicated form, mixing up and down degrees of freedom. We define $\phi_\sigma = \phi_{R,\sigma} + \phi_{L,\sigma}$ and introduce linear combinations of ϕ_σ to describe the charge and spin degrees of freedom,

$$\phi_c = \frac{1}{\sqrt{2}}(\phi_\uparrow + \phi_\downarrow), \quad \phi_s = \frac{1}{\sqrt{2}}(\phi_\uparrow - \phi_\downarrow), \quad (6)$$

Then, we can rewrite $H = \int_0^L dx \mathcal{H}$ as (at $h = 0$):

$$\mathcal{H} = \mathcal{H}_{ph} + \mathcal{H}_{free} + \mathcal{V}_{eff} \quad (7)$$

where

$$\begin{aligned} \mathcal{H}_{ph} &= \frac{1}{2}(\frac{1}{2}\tilde{\delta}_{P,t}^2 + \tilde{\delta}_{P,d}^2) + \frac{1}{2}(\frac{1}{2}\tilde{\delta}_{H,t}^2 + \tilde{\delta}_{H,d}^2) \\ \mathcal{H}_{free} &= \frac{v_c}{4\pi} \left[\frac{1}{K_c}(\partial_x \phi_c)^2 + K_\rho(\partial_x \theta_c)^2 \right] \\ &\quad + \frac{v_s}{4\pi} \left[\frac{1}{K_s}(\partial_x \phi_s)^2 + K_s(\partial_x \theta_s)^2 \right] \\ \mathcal{V}_{eff} &= \mathcal{V}_c + \mathcal{V}_s + \mathcal{V}_{int} \end{aligned} \quad (8)$$

$$\begin{aligned}
\mathcal{V}_c &= \tilde{g}_{1/4} \cos 4\phi_c + \lambda_1 \tilde{g}_P \tilde{\delta}_{P,d} \sin 2\phi_c + \lambda_2 \tilde{g}_H \tilde{\delta}_{H,d} \cos 2\phi_c + \lambda_3 \tilde{g}_V \tilde{\delta}_{P,d} \cos 2\phi_c \\
\mathcal{V}_\sigma &= \tilde{g}_s \cos 2\phi_s \\
\mathcal{V}_{int} &= -\tilde{g}_P \tilde{\delta}_{P,t} \cos(\phi_c - \xi) \cos \phi_s - \tilde{g}_H \tilde{\delta}_{H,t} \sin(\phi_c + \zeta) \cos \phi_s - \lambda_4 \tilde{g}_V \tilde{\delta}_{P,d} \cos 2\phi_c \cos 2\phi_s \\
&\quad + \lambda_5 \tilde{g}_V \tilde{\delta}_{P,t} \cos \phi_s \sin\left(3\phi_c + \xi - \frac{\pi}{4}\right) - \lambda_6 \tilde{g}_V \tilde{\delta}_{P,t} \cos \phi_s \sin\left(\phi_c - \xi + \frac{\pi}{4}\right)
\end{aligned} \tag{9}$$

plus several vertex operators that are kept only when they are commensurate (non-oscillating in space) and constitute relevant perturbations to the Gaussian conformal field theory. In Eqs. (8) and (9) ϕ_c and ϕ_s are the charge and spin fields respectively, $\theta_{c,s}$ are the dual fields defined by $\partial_x \theta_{c,s} = \partial_t \phi_{c,s}$ and the phases ξ and ζ determine the spatial patterns of the tetramerization (see Eq. (4)). The parameters v_c and v_s are the velocity of the charge and spin excitations, and K_c and K_s are the corresponding Tomonaga-Luttinger parameters. From the perturbative calculation presented in²⁷ the coupling constants are given by $\tilde{g}_{1/4} \propto U^2(U - 4V)$ and $\tilde{g}_s \propto U - \alpha U(U - 2V)$ where α is some numerical constant. Besides,

$$\begin{aligned}
\tilde{g}_{1/4} &\propto U^2(U - 4V) \\
\tilde{g}_s &\propto U - \alpha U(U - 2V) \\
\lambda_1 &\propto U \\
\lambda_2 &\propto U \\
\lambda_3 &\propto (U - cte)V \\
\lambda_4 &\propto V \\
\lambda_5 &\propto UV \\
\lambda_6 &\propto (U - cte')V.
\end{aligned} \tag{10}$$

The Eq. (9) is the bosonic self-interaction potential defining a Multi-sine Gordon theory. Extensive analysis of the competition between different harmonics in multi-frequency sine-Gordon theories have been performed in^{28,29,30}, mainly focused on the double sine-Gordon model. The three-frequency case has also been recently discussed in^{31,32,33}. For our purposes it will be enough to perform a semiclassical treatment, as detailed in the next Section.

In order to perform the semiclassical analysis of the ground state and present a schematic phase diagram, we use the bare coefficients in Eq. (9) and we assume the qualitative phenomenological dependence of the coupling constants $\tilde{g}_{1/4}$, $\tilde{g}_s$ and λ 's on the microscopic parameters as in 10.

The aim of the present work is to search for the possibility of elastic deformations that lower the energy with respect to the homogeneous non-deformed case. The simplest analysis of the effective theory Eq. (7), (8) and (9), which has proven to be useful in related cases^{32,33,34,35}, consists in treating the self-interaction terms in Eq. (9) as a classical potential to be evaluated in constant field configurations. Within this approximation the energy

per site depends on four configuration parameters,

$$\epsilon(\phi_c, \phi_s, \xi, \zeta) = \mathcal{V}_{eff}(\phi_c, \phi_s, \xi, \zeta) \tag{11}$$

so that the minima can be found analytically. In the previous expression, ϕ_c and ϕ_s are the charge and spin fields, and the phases ξ and ζ determine the spatial patterns of the tetramerization.

Our results on coupled Holstein-Peierls-Hubbard chain are summarized in Fig. 3. First, we consider the case $\tilde{g}_V = 0$. From the perturbative calculation, the coupling $\tilde{g}_{1/4}$ is positive for small V but becomes negative for large V . When $\tilde{g}_{1/4} > 0$, the phase variable ϕ_c is to be fixed at $\phi_c = \pi/4$, while for $\tilde{g}_{1/4} < 0$, $\phi_c = 0$.

Let us consider finite values of $\tilde{g}_P$ and $\tilde{g}_H$. In Fig. 3(a) we have set $\tilde{g}_{1/4} > 0$ (or $U > 4V$), and we increase $\tilde{g}_P$ from zero (for fixed small $\tilde{g}_H$). Unlike what has been found numerically in¹⁹ (where no distortion patterns were favored, probably due to finite size effects), the phases (ϕ_c, ϕ_s) are locked in $(\frac{\pi}{4}, \frac{\pi}{2})$ (independently of ξ, ζ). In this case, the ground state becomes the $4k_F$ BOW while the site charge density remains uniform ($\langle n_i \rangle = 1/2$), the bond-order is now inhomogeneous and has the form shown in Fig. 1(a). For $\tilde{g}_P$ larger than a critical values $\tilde{g}_P \gtrsim \tilde{g}_P^c(\tilde{g}_H)$, we have $(\xi, \zeta, \phi_c, \phi_s) = (\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{4}, \pi)$ and the BCDW depicted in Fig. 1(b) becomes the ground state. Hence the bond-order pattern now is $t(1 + \alpha)$, $t(1 - \alpha')$, $t(1 + \alpha)$, $t(1 - \alpha')$ and the charge order follows the $\circ - \bullet - \bullet - \circ$ pattern. Now consider the situation when $\tilde{g}_H$ reaches a critical value $\tilde{g}_H^c(\tilde{g}_P)$. In that case, $(\phi_c, \phi_s) = (0, \frac{\pi}{2})$ (independently of ξ, ζ) and the $4k_F$ CDW with $\bullet - \circ - \bullet - \circ$ charge order and uniform bond order becomes the ground state Fig. 1(c). The SP distortion, which corresponds to the $4k_F$ CDW- $2k_F$ SP state (shown in Fig. 1(d) occurs only for large values of $\tilde{g}_H$ and $\tilde{g}_P \approx 0.8$. In this case $(\xi, \zeta, \phi_\rho, \phi_\sigma) = (\frac{\pi}{2}, \pi, 0, \pi)$.

In Fig. 3(b) we can see several important trends when $\tilde{g}_{1/4} < 0$ (or $V > U/4$). First, the size of the $4k_F$ BOW region decreases considerably with decreasing $\tilde{g}_{1/4} < 0$, showing that it takes stronger e-ph coupling to overcome the “natural” tendency towards $\bullet - \circ - \bullet - \circ$ charge order for small $\tilde{g}_P$. For the same reason, the sizes of the $4k_F$ CDW and $4k_F$ CDW- $2k_F$ SP regions increase considerably when $\tilde{g}_{1/4}$ is negative. Next we examine the effect of $\tilde{g}_V$. When $\tilde{g}_V = 0.2$ and $\tilde{g}_{1/4} < 0$, we can see that the $4k_F$ BOW region “disappears” and for $\tilde{g}_P \lesssim 1.5$ the new phases $4k_F$ CDW- $4k_F$ SP and $4k_F$ CDW- $4k_F$ SP + $2k_F$ CDW- $2k_F$ SP dominate. Unlike the previous case with $\tilde{g}_V = 0$, in these phases, there




FIG. 2: Semiclassical energies for the minima in terms of $\tilde{g}_P$ for $\tilde{g}_{1/4} = -1.5$, $\tilde{g}_H = 2.2$ and $\tilde{g}_V = 0$.




FIG. 3: Schematic phase diagram in the plane $\tilde{g}_P$ vs $\tilde{g}_H$ for different values of $\tilde{g}_V$. (a) $\tilde{g}_{1/4} = 1$, $\tilde{g}_V = 0$; (b) $\tilde{g}_{1/4} = -3$, $\tilde{g}_V = 0$; (d) $\tilde{g}_{1/4} = -3$, $\tilde{g}_V = 0.2$. The boundaries of the phases result from level crossings as shown in the Fig. (2). The case with $\tilde{g}_{1/4} > 0$ and $\tilde{g}_V > 0$ is similar to the case (a) and we don't show here. The effect of the Peierls deformation in the nearest-neighbor Coulomb repulsion is more important for large V , that is, $\tilde{g}_{1/4} < 0$.

is now a small dimerization in the bonds (Fig. 1(e) and Fig. 1(f)).

IV. SUMMARY AND DISCUSSIONS

We have investigated the electronic ground-state phase diagram of the one-dimensional Hubbard model at quarter filling with on-site (U) and nearest-neighbor (V) Coulomb repulsion coupled to adiabatic phonons that includes Holstein and Peierls interactions. In addition to the conventional terms, we have included the effects of the Peierls deformation in the nearest-neighbor Coulomb interaction V that may be important when the Coulomb interaction between two sites r_i and r_j is exponentially short ranged, *i.e.* $\exp[-\alpha|r_i - r_j|]/r^\xi$ ($\alpha, \xi > 0$). We have used Abelian bosonization to analyze the low energy properties of the model and we have performed a semiclassical analysis of the bosonized effective theory to identify the possible phases of the ground state. We have found that several charge orders combined with lattice deformations describe the ground states of the system depending on the microscopic parameters U, V and the spin-phonon couplings $\tilde{g}_P, \tilde{g}_H$ and $\tilde{g}_V$. In each of these phases a non trivial elastic deformation is favored, grouping together blocks of two or four sites, while the charge sector adopts different charge orders. A detailed analysis for particular values of $\tilde{g}_V$, shows the following phases at zero temperature:

- For $U > V$ ($\tilde{g}_{1/4} > 0$), $\tilde{g}_V = 0$ and for small and nonzero $\tilde{g}_P$ and $\tilde{g}_H$, the ground state has a dimerized $4k_F$ Bond-Order-wave (BOW) with uniform site charges Fig. 1(a). If $\tilde{g}_P$ is sufficiently large ($\tilde{g}_H$ fixed), there is a second dimerization of the dimerized lattice, leading to the BCDW state (a superposition of the $2k_F$ and $4k_F$ BOWs), accompanied by the $\bullet - \bullet - o - o$ charge order Fig. 1(b). On the other hand, at large $\tilde{g}_H$ and small $\tilde{g}_P$, a $4k_F$ CDW is found, with uniform bonds Fig. 1(c). If we increase $\tilde{g}_P$, the ground state becomes the $4k_F$ CDW- $2k_F$ SP, where the charge order $\bullet - \bullet - o - o$, is accompanied by a $2k_F$ lattice distortion Fig. 1(d).
- For $V > U$ ($\tilde{g}_{1/4} < 0$), the $4k_F$ BOW phase is diminished due to the strong Coulomb repulsion V and phases with charge-order dominate the phase diagram Fig. (3).
- For $V > U$ ($\tilde{g}_{1/4} < 0$), $\tilde{g}_V > 0$, the $4k_F$ BOW disappears because the Coulomb interaction between nearest-neighbor is increased due the Peierls deformation. The CDW's phases has a small contribution of $4k_F$ BOWs combined with the original $2k_F$, giving rise to two new phases: $4k_F$ CDW- $4k_F$ SP and $4k_F$ CDW- $4k_F$ SP + $2k_F$ CDW- $2k_F$ SP

In conclusion, we have argued that the coexistence of charge order and lattice distortions is dominated by $4k_F$

BOW, BCDW for $\tilde{g}_{1/4} > 0$. On the one hand, when $\tilde{g}_{1/4} < 0$, the $4k_F$ CDW and $4k_F$ CDW- $2k_F$ SP dominate if $\tilde{g}_P < \tilde{g}_P^c$, Fig. 3(b); for $\tilde{g}_P > \tilde{g}_P^c$ the BCDW phase dominate. On the other hand, when the phonon-lattice coupling, $\tilde{g}_V$ is taken into account (only for $\tilde{g}_{1/4} < 0$ because is necessary $V > U$), two new phases that we named $4k_F$ CDW- $4k_F$ SP and $4k_F$ CDW- $4k_F$ SP+ $2k_F$ CDW- $2k_F$ SP (Fig. 1(e) and Fig. 1(f) respectively) dominate the phase diagram. These phases differ from the original $4k_F$ CDW and $4k_F$ CDW-SP, in a small dimerization in the bonds, $\tilde{\delta}_{P,d}$ and a small contribution of $2k_F$ CDW in the last one.

Numerical analysis of the present system including Peierls deformations in the nearest-neighbor coulomb interaction would be helpful in order to determine the phase boundaries quantitatively. We hope that our predictions will stimulate further X-ray or neutron scattering experiments on compounds like Bechgaard salts. They should be able to distinguish the different distortion

patters predicted here. Since the effects of the Peierls deformations in the nearest-neighbor coulomb interaction is proportional to V (see Eq. (1)), we expect that the effects of this term will be more important in compounds where V is larger.

V. ACKNOWLEDGMENTS

We thank A. Honecker, G.I. Japaridze, C.A. Lamas, M. Moliner, D. Poilblanc, P. Pujol and G.L. Rossini for helpful discussions.

This work was partially supported by the ESF grant INSTANS, ECOS-Sud Argentina-France collaboration (Grant No A04E03), PICS CNRS-Conicet (Grant No. 18294), PICT ANCYPT (Grant No 20350), and PIP CONICET (Grant No 5037).

-
- ¹ For reviews on Bechgaard salts see *e.g.* D. Jérôme and H.J. Schulz, Adv. Phys. **31**, 299 (1982); D. Jerome, *Organic superconductors: From (TMTSF)₂PF₆ to fullerenes* (Marcel Dekker, New York, 1994), 405. T. Ishiguro and K. Yamaji, Springer Series Solid State Vol. **88**, Springer-Verlag, Berlin (1990). T. Ishiguro, K. Yamaji and G. Saito, *Organics Superconductors*, Springer-Verlag, New York, (1998).
 - ² K. Hiraki and K. Kanoda, Phys. Rev. Lett. **80**, 4737 (1998).
 - ³ S. Mazumdar, D. Campbell, R.T. Clay and S. Ramasesha, Phys. Rev. Lett. **82**, 2411 (1999).
 - ⁴ K. Hiraki and K. Kanoda, Phys. Rev. Lett. **82**, 2412 (1999).
 - ⁵ N. Biskup, J.A.A.J. Perenboom, J.S. Brooks, and J.S. Qualls, Solid State Comm. **107**, 503 (1998).
 - ⁶ D.S. Chow, F. Zamborsky, B. Alavi, D.J. Tantillo, A. Baur, C.A. Merlic, and S.E. Brown, Phys. Rev. Lett. **85**, 1698 (2000).
 - ⁷ F. Nad, P. Monceau, C. Carcel and J.M. Fabre, J. Phys. Cond. Matt. **12**, L435 (2000).
 - ⁸ F. Nad, P. Monceau, C. Carcel, and J.M. Fabre, Phys. Rev. B **62**, 1753 (2000).
 - ⁹ P. Monceau, F.Y. Nad, and S. Brazovskii, Phys. Rev. Lett. **86**, 4080 (2001).
 - ¹⁰ K. Miyagawa, A. Kawamoto and K. Kanoda, Phys. Rev. B **62**, R7679 (2000).
 - ¹¹ K.C. Ung, S. Mazumdar and D. Toussaint, Phys. Rev. Lett. **73**, 2603 (1994).
 - ¹² S. Mazumdar, S. Ramasesha, R.T. Clay and D.K. Campbell, Phys. Rev. Lett. **82**, 1522 (1999).
 - ¹³ N. Kobayashi and M. Ogata, J. Phys. Soc. Jpn. **66**, 3356 (1997).
 - ¹⁴ N. Kobayashi, M. Ogata and K. Yonemitsu, J. Phys. Soc. Jpn. **67**, 1098 (1998).
 - ¹⁵ J. Riera and D. Poilblanc, Phys. Rev. B **59**, 2667 (1999).
 - ¹⁶ J. Riera and D. Poilblanc, Phys. Rev. B **62**, R16243 (2000).
 - ¹⁷ H. Seo and H. Fukuyama, J. Phys. Soc. Jpn. **66**, 1249 (1997).
 - ¹⁸ H. Seo, J. Phys. Soc. Jpn. **69**, 805 (2000).
 - ¹⁹ R.T. Clay, S. Mazumdar and D.K. Campbell, Phys. Rev. B **67**, 115121 (2003).
 - ²⁰ D.C. Cabra, A. De Martino, A. Honecker, P. Pujol, and P. Simon, Phys. Rev. B **63**, 094406 (2001).
 - ²¹ For a recent review on the Hubbard chain coupled to adiabatic phonons, see for example chapter IV in http://eprints-scd-ulp.u-strasbg.fr:8080/1100/01/MOLINER_Marion.2
 - ²² K. Penc and F. Mila, Phys. Rev. B **49**, 9670 (1994).
 - ²³ H.Q. Lin, *et al*, Proceedings of the 1993 NATO ARW on *The Physics and Mathematical Physics of the Hubbard Model*, edited by D. Baeriswyl *et al*, Plenum, New York (1995).
 - ²⁴ T. Giamarchi, *Quantum Physics in One Dimension*, Oxford University Press, Oxford (2004).
 - ²⁵ See for instance D.C. Cabra and P. Pujol, in *Quantum Magnetism*, Lect. Notes in Phys. **645** 253 (2004).
 - ²⁶ D. Sènèchal, *An introduction to bosonization*, arXiv: cond-mat 9908262v1 (1999) or in the book D. Sènèchal, A.-M. Tremblay and C. Bourbonnais *Theoretical Methods for Strongly Correlated Electrons*, Springer, 1 edition (October 1, 2003).
 - ²⁷ H. Yoshioka, M. Tsuchiizu and Y. Suzumura, J. Phys. Soc. Jap. **69**, 651 (2000).
 - ²⁸ G. Delfino and G. Mussardo, Nucl. Phys. B **516**, 675 (1998).
 - ²⁹ M. Fabrizio, A.O. Gogolin, and A.A. Nersesyan, Nucl. Phys. B **580**, 647 (2000).
 - ³⁰ Z. Bajnok, L. Palla, G. Takacs, and F. Wagner, Nucl. Phys. B **601**, 503 (2001).
 - ³¹ G.Z. Tóth, J. Phys. A **37**, 9631 (2004).
 - ³² H.D. Rosales, D.C. Cabra, M.D. Grynberg, G.L. Rossini, and T. Vekua, Phys. Rev. B **75**, 174446 (2007).
 - ³³ H.D. Rosales and G.L. Rossini, Phys. Rev. B **76**, 224404 (2007).
 - ³⁴ P. Lecheminant and E. Orignac, Phys. Rev. B **69**, 174409 (2004).
 - ³⁵ K. Hida and I. Affleck, J. Phys. Soc. Jpn. **74**, 1849 (2005).